Integral Transforms and Special Functions
Vol. 00, No. 00, January 2009, 1–20

arXiv:1202.1811v1 [math.CA] 8 Feb 2012

RESEARCH ARTICLE

Fourier expansions for a logarithmic fundamental solution of the polyharmonic equation

Howard S. Cohl[a,b*]

[a]*Applied and Computational Mathematics Division, Information Technology Laboratory, National Institute of Standards and Technology, Gaithersburg, Maryland, U.S.A.* ;
[b]*Department of Mathematics, University of Auckland, Auckland, New Zealand*

(*v3.5 released October 2008*)

In even-dimensional Euclidean space for integer powers of the Laplacian greater than or equal to the dimension divided by two, a fundamental solution for the polyharmonic equation has logarithmic behavior. We give two approaches for developing a Fourier expansion of this logarithmic fundamental solution. The first approach is algebraic and relies upon the construction of two-parameter polynomials. We describe some of the properties of these polynomials, and use them to derive the Fourier expansion for a logarithmic fundamental solution of the polyharmonic equation. The second approach depends on the computation of parameter derivatives of Fourier series for a power-law fundamental solution of the polyharmonic equation. The resulting Fourier series is given in terms of sums over associated Legendre functions of the first kind. We conclude by comparing the two approaches and giving the azimuthal Fourier series for a logarithmic fundamental solution of the polyharmonic equation in rotationally-invariant coordinate systems.

Keywords: fundamental solutions; polyharmonic equation; Fourier series; polynomials; associated Legendre functions

AMS Subject Classification: 35A08; 31B30; 31C12; 33C05; 42A16

1. Introduction

Solutions of the polyharmonic equation (powers of the Laplacian operator) are ubiquitous in many areas of computational, pure, applied mathematics, physics and engineering. We concern ourselves, in this paper, with a fundamental solution of the polyharmonic equation (Laplace, biharmonic, etc.), which by convolution yields a solution to the inhomogeneous polyharmonic equation. Solutions to inhomogeneous polyharmonic equations are useful in many physical applications including those areas related to Poisson's equation such as Newtonian gravity, electrostatics, magnetostatics, quantum direct and exchange interactions (cf. §1 in Cohl & Dominici (2010) [4]), etc. Furthermore, applications of higher-powers of the Laplacian include such varied areas as minimal surfaces [13], Continuum Mechanics [9], Mesh deformation [7], Elasticity [10], Stokes Flow [8], Geometric Design [20], Cubature formulae [17], mean value theorems (cf. Pizzetti's formula) [14], and Hartree-Fock calculations of nuclei [21].

*Corresponding author. Email: howard.cohl@nist.gov

ISSN: 1065-2469 print/ISSN 1476-8291 online
© 2009 Taylor & Francis

It is a well-known fact (see for instance Schwartz (1950) ([16], p. 45), Gel′fand & Shilov (1964) ([6], p. 202) that a fundamental solution of the polyharmonic equation on d-dimensional Euclidean space is given by combinations of power-law and logarithmic functions of the global distance between two points. In a recent paper (Cohl & Dominici (2010) [4]), we derived a complex identity which determined the Fourier coefficients of a power-law fundamental solution of the polyharmonic equation. The Fourier coefficients were seen to be given in terms of associated Legendre functions. The present work is concerned with computing the Fourier coefficients of a logarithmic fundamental solution of the polyharmonic equation. One obtains a logarithmic fundamental solution for the polyharmonic equation only in even-dimensional Euclidean space and only when the power of the Laplacian is greater than or equal to the dimension divided by two. The most familiar example of a logarithmic fundamental solution of the polyharmonic equation occurs in two-dimensions, for a single-power of the Laplacian, i.e., Laplace's equation.

We present two different approaches for obtaining Fourier series of a logarithmic fundamental solution for the polyharmonic equation. The first approach is algebraic and involves the generation of a certain set of naturally arising two-index polynomials which we refer to as logarithmic polynomials. The second approach starts with the main result from Cohl & Dominici (2010) [4] and determines the Fourier series expansion for a logarithmic fundamental solution of the polyharmonic equation through parameter differentiation. Series expansions for fundamental solutions of linear partial differential equations such as the polyharmonic equation are extremely useful in determining Dirichlet boundary values for solutions on interior domains (see for example Cohl & Tohline (1999) [5]).

This paper is organized as follows. In §2 we introduce the problem. In §3 we describe our algebraic approach to computing a Fourier series of a logarithmic fundamental solution of the polyharmonic equation. In §4 we give our limit derivative approach for computing the Fourier series of a logarithmic fundamental solution of the polyharmonic equation. In §5 we give some comparisons between the two approaches. In §6 we use the results presented in the previous sections to obtain azimuthal Fourier expansions for a logarithmic fundamental solution of the polyharmonic equation in rotationally-invariant coordinate systems which parametrize points in d-dimensional Euclidean space. In Appendix A we present some necessary formulae relating to differentiation of associated Legendre functions of the first kind with respect to the degree. In Appendix B we present some of the properties of the logarithmic polynomials.

Throughout this paper we rely on the following definitions. The set of natural numbers is given by $\mathbf{N} := \{1, 2, 3, \ldots\}$, the set $\mathbf{N}_0 := \{0, 1, 2, \ldots\} = \mathbf{N} \cup \{0\}$, the set of integers is given by $\mathbf{Z} := \{0, \pm 1, \pm 2, \ldots\}$, and the set $\mathbf{Q}$ represents the rational numbers. For $a_1, a_2, \ldots \in \mathbf{C}$, if $i, j \in \mathbf{Z}$ and $j < i$ then $\sum_{n=i}^{j} a_n = 0$ and $\prod_{n=i}^{j} a_n = 1$, where $\mathbf{C}$ represents the complex numbers. The set $\mathbf{R}$ represents the real numbers.

2. Fundamental solution of the polyharmonic equation and the non-logarithmic Fourier series

If Φ satisfies the polyharmonic equation given by

$$(-\Delta)^k \Phi(\mathbf{x}) = 0, \tag{1}$$

where $\mathbf{x} \in \mathbf{R}^d$, $\Delta : C^p(\mathbf{R}^d) \to C^{p-2}(\mathbf{R}^d)$, for $p \geq 2$ is the Laplacian operator defined by $\Delta := \frac{\partial^2}{\partial x_1^2} + \ldots + \frac{\partial^2}{\partial x_d^2}$, $k \in \mathbf{N}$ and $\Phi \in C^{2k}(\mathbf{R}^d)$, then Φ is called polyharmonic. If the power k of the Laplacian equals two, then (1) is called the biharmonic equation and Φ is called biharmonic. The inhomogeneous polyharmonic equation is given by

$$(-\Delta)^k \Phi(\mathbf{x}) = \rho(\mathbf{x}), \tag{2}$$

where we take ρ to be an integrable function so that a solution to (2) exists. A fundamental solution for the polyharmonic equation on $\mathbf{R}^d$ is a function $\mathfrak{g}_k^d : (\mathbf{R}^d \times \mathbf{R}^d) \setminus \{(\mathbf{x}, \mathbf{x}) : \mathbf{x} \in \mathbf{R}^d\} \to \mathbf{R}$ which satisfies the equation

$$(-\Delta)^k \mathfrak{g}_k^d(\mathbf{x}, \mathbf{x}') = c\delta(\mathbf{x} - \mathbf{x}'), \tag{3}$$

for some $c \in \mathbf{R}, c \neq 0$, where δ is the Dirac delta function and $\mathbf{x}' \in \mathbf{R}^d$. When $c = 1$, we call a fundamental solution of the polyharmonic equation normalized, and denote it by $\mathcal{G}_k^d : (\mathbf{R}^d \times \mathbf{R}^d) \setminus \{(\mathbf{x}, \mathbf{x}) : \mathbf{x} \in \mathbf{R}^d\} \to \mathbf{R}$. The Euclidean inner product $(\cdot, \cdot) : \mathbf{R}^d \times \mathbf{R}^d \to \mathbf{R}$ defined by $(\mathbf{x}, \mathbf{x}') := x_1 x_1' + \ldots + x_d x_d'$, induces a norm (the Euclidean norm) $\|\cdot\| : \mathbf{R}^d \to [0, \infty)$, on the finite-dimensional vector space $\mathbf{R}^d$, given by $\|\mathbf{x}\| := \sqrt{(\mathbf{x}, \mathbf{x})}$. In the rest of this paper, we will use the gamma function $\Gamma : \mathbf{C} \setminus -\mathbf{N}_0 \to \mathbf{C}$, which is a natural generalization of the factorial function (see for instance Chapter 5 in Olver *et al.* (2010) [15]). A fundamental solution of the polyharmonic equation is given by the following theorem.

THEOREM 2.1 *Let $d, k \in \mathbf{N}$. Define*

$$\mathcal{G}_k^d(\mathbf{x}, \mathbf{x}') = \begin{cases} \dfrac{(-1)^{k+d/2+1} \ \|\mathbf{x} - \mathbf{x}'\|^{2k-d}}{(k-1)! \ (k-d/2)! \ 2^{2k-1}\pi^{d/2}} \left(\log \|\mathbf{x} - \mathbf{x}'\| - \beta_{k-d/2,d}\right) & \\ & \text{if } d \text{ even}, \ k \geq d/2, \\ \dfrac{\Gamma(d/2-k)\|\mathbf{x} - \mathbf{x}'\|^{2k-d}}{(k-1)! \ 2^{2k}\pi^{d/2}} & \text{otherwise,} \end{cases}$$

where $\beta_{p,d} \in \mathbf{Q}$ is defined as $\beta_{p,d} := \frac{1}{2}\left[H_p + H_{d/2+p-1} - H_{d/2-1}\right]$, with H_j being the jth harmonic number

$$H_j := \sum_{i=1}^{j} \frac{1}{i},$$

then $\mathcal{G}_k^d$ is a normalized fundamental solution for $(-\Delta)^k$ on Euclidean space $\mathbf{R}^d$.

Proof. See Cohl (2010) [3] and Boyling (1996) [2].

A separable rotationally-invariant coordinate system for the polyharmonic equation (1) on $\mathbf{R}^d$ is given by

$$\left.\begin{aligned} x_1 &= R(\xi_1, \ldots, \xi_{d-1}) \cos\phi \\ x_2 &= R(\xi_1, \ldots, \xi_{d-1}) \sin\phi \\ x_3 &= x_3(\xi_1, \ldots, \xi_{d-1}) \\ &\vdots \\ x_d &= x_d(\xi_1, \ldots, \xi_{d-1}) \end{aligned}\right\}, \tag{4}$$

which is described by an angle $\phi \in \mathbf{R}$ and $(d-1)$-curvilinear coordinates $(\xi_1, \ldots, \xi_{d-1})$. A separable rotationally-invariant coordinate system transforms the polyharmonic equation into a set of d-uncoupled ordinary differential equations with separation constants $m \in \mathbf{Z}$ and k_j for $1 \leq j \leq d-2$. For a separable rotationally-invariant coordinate system, this uncoupling is accomplished, in general, by assuming a solution to (1) of the form

$$\Phi(x) = e^{im\phi}\, \mathcal{R}(\xi_1, \ldots, \xi_{d-1}) \prod_{i=1}^{d-1} A_i(\xi_i, m, k_1, \ldots, k_{d-2}),$$

where the domains of the functions $\mathcal{R}$ and A_i, for $1 \leq i \leq d-1$, and the constants k_j for $1 \leq j \leq d-1$, depend on the specific rotationally-invariant coordinate system. A rotationally-invariant coordinate system parametrizes points on the $(d-1)$-dimensional half-hyperplane given by $\phi = const.$ and $R \geq 0$ using curvilinear coordinates $(\xi_1, \ldots, \xi_{d-1})$. (For a general description of the theory of separation of variables see Miller (1977) [12].) The Euclidean distance between two points $\mathbf{x}, \mathbf{x}' \in \mathbf{R}^d$, expressed in a rotationally-invariant coordinate system, is given by

$$\|\mathbf{x} - \mathbf{x}'\| = \sqrt{2RR'}\left[\chi - \cos(\phi - \phi')\right]^{1/2},$$

where the toroidal parameter $\chi > 1$, is given by

$$\chi := \frac{R + R'^2 + \sum_{i=3}^{d} (x_i - x_i')^2}{2RR'}, \tag{5}$$

where $R, R' \in [0, \infty)$ are defined in (4) for $\mathbf{x}, \mathbf{x}' \in \mathbf{R}^d$. The hypersurfaces given by $\chi > 1$ equals constant are independent of coordinate system and represent hyper-tori of revolution.

From Theorem 2.1 we see that, apart from multiplicative constants, the algebraic expression $\mathfrak{l}_k^d : (\mathbf{R}^d \times \mathbf{R}^d) \setminus \{(\mathbf{x}, \mathbf{x}) : \mathbf{x} \in \mathbf{R}^d\} \to \mathbf{R}$ of an unnormalized fundamental solution for the polyharmonic equation in Euclidean space $\mathbf{R}^d$ for d even, $k \geq d/2$, is given by

$$\mathfrak{l}_k^d(\mathbf{x}, \mathbf{x}') := \|\mathbf{x} - \mathbf{x}'\|^{2k-d} \left(\log \|\mathbf{x} - \mathbf{x}'\| - \beta_{k-d/2,d}\right). \tag{6}$$

By expressing $\mathfrak{l}_k^d$ in a rotationally-invariant coordinate system (4) we obtain

$$\begin{aligned}\mathfrak{l}_k^d(\mathbf{x}, \mathbf{x}') = \left(2RR'\right)^p \left[\frac{1}{2}\log\left(2RR'\right) - \beta_{p,d}\right] \left[\chi - \cos(\phi - \phi')\right]^p \\ + \frac{1}{2}\left(2RR'\right)^p \left[\chi - \cos(\phi - \phi')\right]^p \log\left[\chi - \cos(\phi - \phi')\right],\end{aligned} \tag{7}$$

where $p = k - d/2 \in \mathbf{N}_0$. For the polyharmonic equation in even-dimensional Euclidean space $\mathbf{R}^d$ with $1 \leq k \leq d/2 - 1$, apart from multiplicative constants, the algebraic expression for an unnormalized fundamental solution of the polyharmonic equation $\mathfrak{h}_k^d : (\mathbf{R}^d \times \mathbf{R}^d) \setminus \{(\mathbf{x}, \mathbf{x}) : \mathbf{x} \in \mathbf{R}^d\} \to \mathbf{R}$ is given by

$$\mathfrak{h}_k^d(\mathbf{x}, \mathbf{x}') := \|\mathbf{x} - \mathbf{x}'\|^{2k-d}.$$

By expressing $\mathfrak{h}_k^d$ in a rotationally-invariant coordinate system we obtain

$$\mathfrak{h}_k^d(\mathbf{x}, \mathbf{x}') = \left(2RR'\right)^{-q} \left[\chi - \cos(\phi - \phi')\right]^{-q}, \tag{8}$$

where $q = 2k - d$.

By examining (7) and (8), we see that for computation of Fourier expansions about the azimuthal separation angle $(\phi - \phi')$ of $\mathfrak{l}_k^d$ and $\mathfrak{h}_k^d$, all that is required is to compute the Fourier cosine series for the following three functions $f_\chi, h_\chi : \mathbf{R} \to (0, \infty)$ and $g_\chi : \mathbf{R} \to \mathbf{R}$ defined as

$$f_\chi(\psi) := (\chi - \cos\psi)^p,$$

$$g_\chi(\psi) := (\chi - \cos\psi)^p \log(\chi - \cos\psi), \qquad \text{and}$$

$$h_\chi(\psi) := (\chi - \cos\psi)^{-q},$$

where $p \in \mathbf{N}_0$, $q \in \mathbf{N}$ and $\chi > 1$ is a fixed parameter.

The Fourier series of f_χ is given in Cohl & Dominici (2010) [4] (cf. (4.4) therein), namely

$$(z - \cos\psi)^p = (z^2 - 1)^{p/2} \sum_{n=0}^{p} \epsilon_n \cos(n\psi) \frac{(-p)_n (p-n)!}{(p+n)!} P_p^n \left(\frac{z}{\sqrt{z^2-1}}\right), \tag{9}$$

where the Neumann factor $\epsilon_n = 2 - \delta_{n,0}$ commonly occurs in Fourier series, $\delta_{n,0}$ is the Kronecker delta, and

$$(z)_n := \prod_{i=1}^{n} (z + i - 1),$$

for $z \in \mathbf{C}$ and $n \in \mathbf{N}_0$, is the Pochhammer symbol (rising factorial). We have used Whipple's formula in (9) (see for instance, (8.2.7) in Abramowitz & Stegun (1972) [1]) to convert the associated Legendre function of the second kind $Q_\nu^\mu : (1, \infty) \to \mathbf{C}$ appearing in [4] to the associated Legendre function of the first kind $P_\nu^\mu : (1, \infty) \to \mathbf{R}$. The associated Legendre function of the first kind can be defined using the Gauss hypergeometric function, namely (Magnus, Oberhettinger & Soni (1966) [11], p. 153)

$$P_\nu^\mu(z) := \frac{1}{\Gamma(1-\mu)} \left(\frac{z+1}{z-1}\right)^{\mu/2} {}_2F_1\left(-\nu, \nu+1; 1-\mu; \frac{1-z}{2}\right).$$

The Gauss hypergeometric function ${}_2F_1 : \mathbf{C} \times \mathbf{C} \times (\mathbf{C} \backslash -\mathbf{N}_0) \times \{z \in \mathbf{C} : |z| < 1\} \to \mathbf{C}$ can be defined in terms of the following infinite series

$${}_2F_1(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{n!(c)_n} z^n$$

(see for instance Chapter 15 in Olver *et al.* (2010) [15]).

The Fourier series of h_χ is given in Cohl & Dominici (2010) [4] (Whipple formula

(8.2.7) in Abramowitz & Stegun (1972) [1] and cf. (4.5) therein), namely

$$\frac{1}{(z-\cos\psi)^q} = \frac{(z^2-1)^{-q/2}}{(q-1)!}\sum_{n=0}^{\infty}\epsilon_n\cos(n\psi)(n+q-1)!P_{q-1}^{-n}\left(\frac{z}{\sqrt{z^2-1}}\right), \quad (10)$$

where $q \in \mathbf{N}$. Since the Fourier series of h_χ is computed in Cohl & Dominici (2010) [4], we understand how to compute Fourier expansions of $\mathfrak{h}_k^d$ (8) in separable rotationally-invariant coordinate systems. In order to compute Fourier expansion of $\mathfrak{l}_k^d$ (7) in separable rotationally-invariant coordinate systems, all that remains is to determine the Fourier series of g_χ. This is the goal of the next two sections.

3. Algebraic approach to the logarithmic Fourier series

Since $\chi > 1$, one may make the substitution $\chi = \cosh\eta$ to evaluate the Fourier series of g_χ. For instance, it is given in the form of $(\cosh\eta-\cos\psi)^p\log(\cosh\eta-\cos\psi)$, where $p \in \mathbf{N}_0$. For $p = 0$ the result is well-known (see for instance Magnus, Oberhettinger & Soni (1966) [11], p. 259)

$$\log(\cosh\eta-\cos\psi) = \eta - \log 2 - 2\sum_{n=1}^{\infty}\frac{e^{-n\eta}}{n}\cos(n\psi), \quad (11)$$

which as we will see, should be compared with (10) for $q = 1$, namely

$$\frac{1}{\cosh\eta-\cos\psi} = \frac{1}{\sinh\eta}\sum_{n=0}^{\infty}\epsilon_n\cos(n\psi)e^{-n\eta}. \quad (12)$$

Note that for $\eta > 0$ we may write e^η and therefore η as a function of $\cosh\eta$ since $\sinh\eta = \sqrt{\cosh^2\eta-1}$, $e^\eta = \cosh\eta+\sqrt{\cosh^2\eta-1}$, and therefore $\eta = \log\left(\cosh\eta+\sqrt{\cosh^2\eta-1}\right)$. Now examine the $p = 1$ case for g_χ. If we multiply both sides of (11) by $(\cosh\eta-\cos\psi)$ and take advantage of the formula

$$\cos(n\psi)\cos\psi = \frac{1}{2}\Big\{\cos[(n+1)\psi]+\cos[(n-1)\psi]\Big\}, \quad (13)$$

then we have

$$\begin{aligned}(\cosh\eta-\cos\psi)\log(\cosh\eta-\cos\psi) &= (\eta-\log 2)\cosh\eta \\ &- (\eta-\log 2)\cos\psi - 2\cosh\eta\sum_{n=1}^{\infty}\frac{e^{-n\eta}}{n}\cos(n\psi) \\ &+ \sum_{n=1}^{\infty}\frac{e^{-n\eta}}{n}\cos[(n+1)\psi]+\sum_{n=1}^{\infty}\frac{e^{-n\eta}}{n}\cos[(n-1)\psi]. \quad (14)\end{aligned}$$

Collecting the contributions to the Fourier cosine series, we obtain

$$\begin{aligned}(\cosh\eta-\cos\psi)\log(\cosh\eta-\cos\psi) &= (1+\eta-\log 2)\cosh\eta\\ &\quad -\sinh\eta+\cos\psi\left(\log 2-1-\eta-\frac{1}{2}e^{-2\eta}\right)\\ &\quad +2\sum_{n=2}^{\infty}\frac{e^{-n\eta}\cos n\psi}{n(n^2-1)}(\cosh\eta+n\sinh\eta). \end{aligned}\tag{15}$$

If we compare (15) with (10) for $q=1$, namely

$$\frac{1}{(\cosh\eta-\cos\psi)^2}=\frac{1}{\sinh^3\eta}\sum_{n=0}^{\infty}\epsilon_n\cos(n\psi)e^{-n\eta}(\cosh\eta+n\sinh\eta),\tag{16}$$

we notice that the factor $(\cosh\eta+n\sinh\eta)$ appears in both series.

For $p=2$ in g_χ, we use (13) and similarly have

$$\begin{aligned}&(\cosh\eta-\cos\psi)^2\log(\cosh\eta-\cos\psi)\\ &\quad=(\eta-\log 2)\cosh^2\eta-2(\eta-\log 2)\cosh\eta\cos\psi\\ &\quad+(\eta-\log 2)\cos^2\psi-(2\cosh^2\eta+1)\sum_{n=1}^{\infty}\frac{e^{-n\eta}}{n}\cos(n\psi)\\ &\quad+2\cosh\eta\sum_{n=1}^{\infty}\frac{e^{-n\eta}}{n}\cos[(n+1)\psi]+2\cosh\eta\sum_{n=1}^{\infty}\frac{e^{-n\eta}}{n}\cos[(n-1)\psi]\\ &\quad-\frac{1}{2}\sum_{n=1}^{\infty}\frac{e^{-n\eta}}{n}\cos[(n+2)\psi]-\frac{1}{2}\sum_{n=1}^{\infty}\frac{e^{-n\eta}}{n}\cos[(n-2)\psi].\end{aligned}$$

If we collect the contributions of the Fourier cosine series, we obtain

$$\begin{aligned}&(\cosh\eta-\cos\psi)^2\log(\cosh\eta-\cos\psi)=(\eta-\log 2)\left(\cosh^2\eta+\frac{1}{2}\right)\\ &+2\cosh\eta\,e^{-\eta}-\frac{1}{4}e^{-2\eta}+\left[-2(\eta-\log 2)\cosh\eta-\left(2\cosh^2\eta+\frac{3}{2}\right)e^{-\eta}\right.\\ &\left.+\cosh\eta\,e^{-2\eta}-\frac{1}{6}e^{-3\eta}\right]\cos\psi+\left[\frac{1}{2}(\eta-\log 2)+2\cosh\eta\,e^{-\eta}\right.\\ &\left.-\frac{1}{2}(2\cosh^2\eta+1)e^{-2\eta}+\frac{2}{3}\cosh\eta\,e^{-3\eta}-\frac{1}{8}e^{-4\eta}\right]\cos 2\psi\\ &-4\sum_{n=3}^{\infty}\frac{e^{-n\eta}\cos(n\psi)}{n(n^2-1)(n^2-4)}\left[(n^2-1)\sinh^2\eta+3n\sinh\eta\cosh\eta+3\cosh^2\eta\right].\end{aligned}\tag{17}$$

By comparing (17) with (10) for $q = 2$, namely

$$\frac{1}{(\cosh\eta - \cos\psi)^3} = \frac{1}{2\sinh^5\eta}$$
$$\times \sum_{n=0}^{\infty} \epsilon_n \cos(n\psi) e^{-n\eta} \left[(n^2-1)\sinh^2\eta + 3n\sinh\eta\cosh\eta + 3\cosh^2\eta\right], \quad (18)$$

then we notice that the factor $\left((n^2-1)\sinh^2\eta + 3n\sinh\eta\cosh\eta + 3\cosh^2\eta\right)$ appears in both series. We will demonstrate in §5, why the identification mentioned in (16) and (18) occurs.

This algebraic approach for determining the Fourier series of g_χ will now be generalized. By starting with (11) and repeatedly multiplying by factors of $(\cosh\eta - \cos\psi)$, we see that the general Fourier series of g_χ can be given in terms of a sequence of polynomials $R_p^k : (1,\infty) \to \mathbf{R}$, with $p \in \mathbf{N}_0$ and $k \in \mathbf{Z}$, as

$$(\cosh\eta - \cos\psi)^p \log(\cosh\eta - \cos\psi) = (\eta - \log 2)(\cosh\eta - \cos\psi)^p$$
$$+ 2\sum_{k=-p}^{p} (-1)^{k+1} R_p^k(\cosh\eta) \sum_{n=1}^{\infty} \frac{e^{-n\eta}}{n} \cos[(n+k)\psi]. \quad (19)$$

We will refer to R_p^k as logarithmic polynomials with argument $x = \cosh\eta$ (in our notation p and k are both indices) (See Appendix B for a description of some of the properties of the logarithmic polynomials).

The double sum in (19) is simplified by making the replacement $n + k \mapsto n$. It then follows that the resulting double sum naturally breaks into two disjoint regions, one triangular

$$\mathcal{A} := \{(k,n) : -p \le k \le p-1,\ k+1 \le n \le p\},$$

with $p(2p+1)$ terms and the other infinite rectangular

$$\mathcal{B} := \{(k,n) : -p \le k \le p,\ p+1 < n < \infty\}.$$

By rearranging the order of the k and n summations in (19), we derive

$$(\cosh\eta - \cos\psi)^p \log(\cosh\eta - \cos\psi) = (\eta - \log 2)(\cosh\eta - \cos\psi)^p$$
$$+ \sum_{n=0}^{p} \cos(n\psi) e^{-n\eta}\, \mathfrak{r}_{n,p}^{-p,n-1}(\cosh\eta) + \sum_{n=1}^{p-1} \cos(n\psi) e^{n\eta}\, \mathfrak{r}_{-n,p}^{-p,-n-1}(\cosh\eta)$$
$$+ \sum_{n=p+1}^{\infty} \frac{\cos(n\psi) e^{-n\eta}}{n(n^2-1)\cdots(n^2-p^2)}\, \Re_{n,p}(\cosh\eta), \quad (20)$$

where $\mathfrak{r}_{n,p}^{k_1,k_2}, \Re_{n,p} : (1,\infty) \to \mathbf{R}$ are defined as

$$\mathfrak{r}_{n,p}^{k_1,k_2}(\cosh\eta) := 2\sum_{k=k_1}^{k_2} \frac{(-1)^{k+1} e^{k\eta} R_p^k(\cosh\eta)}{n-k},$$

and

$$\Re_{n,p}(\cosh\eta) := \frac{(n+p)!}{(n-p-1)!}\mathfrak{r}_{n,p}^{-p,p}(\cosh\eta), \quad (n \geq p+1),$$

respectively. We can also write the Fourier series directly in terms of the logarithmic polynomials R_p^k as follows

$$\begin{aligned}(\cosh\eta-\cos\psi)^p\log(\cosh\eta-\cos\psi) &= (\eta-\log 2)(\cosh\eta-\cos\psi)^p\\ &+2\sum_{n=0}^{p}\cos(n\psi)e^{-n\eta}\sum_{k=-p}^{n-1}\frac{(-1)^{k+1}e^{k\eta}R_p^k(\cosh\eta)}{n-k}\\ &-2\sum_{n=1}^{p-1}\cos(n\psi)e^{n\eta}\sum_{k=-p}^{-n-1}\frac{(-1)^{k+1}e^{k\eta}R_p^k(\cosh\eta)}{n+k}\\ &+2\sum_{n=p+1}^{\infty}\cos(n\psi)e^{-n\eta}\sum_{k=-p}^{p}\frac{(-1)^{k+1}e^{k\eta}R_p^k(\cosh\eta)}{n-k}. \end{aligned} \tag{21}$$

Note that by using (9), then we can express $(\cosh\eta-\cos\psi)^p$ as a Fourier series, namely

$$(\cosh\eta-\cos\psi)^p = \sinh^p\eta\sum_{n=0}^{p}\epsilon_n\cos(n\psi)\frac{(-p)_n(p-n)!}{(p+n)!}P_p^n(\coth\eta), \tag{22}$$

where $p\in\mathbf{N}_0$.

If we define the function $\mathfrak{P}_{n,p}:(1,\infty)\to\mathbf{R}$ such that

$$\mathfrak{P}_{n,p}(\cosh\eta) := \begin{cases} \mathfrak{r}_{0,p}^{-p,-1}(\cosh\eta) & \text{if } n=0,\\ D_{n,p}(\cosh\eta)+E_{n,p}(\cosh\eta) & \text{if } 1\leq n\leq p-1,\\ e^{-p\eta}\mathfrak{r}_{p,p}^{-p,p-1}(\cosh\eta) & \text{if } 1\leq n=p,\\ \dfrac{e^{-n\eta}}{(n^2-p^2)\cdots(n^2-1)n}\,\Re_{n,p}(\cosh\eta) & \text{if } n\geq p+1, \end{cases}$$

where $D_{n,p}, E_{n,p}:(1,\infty)\to\mathbf{R}$ are defined as

$$D_{n,p}(\cosh\eta) = \begin{cases} e^{n\eta}\mathfrak{r}_{-n,p}^{-p,-n-1}(\cosh\eta) & \text{if } p\geq 2,\\ 0 & \text{if } p\in\{0,1\}, \end{cases}$$

and

$$E_{n,p}(\cosh\eta) = \begin{cases} e^{-n\eta}\mathfrak{r}_{n,p}^{-p,n-1}(\cosh\eta) & \text{if } p\geq 1,\\ 0 & \text{if } p=0, \end{cases}$$

respectively, then we can write (20) as

$$(\cosh\eta-\cos\psi)^p\log(\cosh\eta-\cos\psi)=(\eta-\log 2)(\cosh\eta-\cos\psi)^p + \sum_{n=0}^{\infty}\cos(n\psi)\mathfrak{P}_{n,p}(\cosh\eta).$$

In fact, if we use (22), then we can express g_χ as

$$(\cosh\eta-\cos\psi)^p\log(\cosh\eta-\cos\psi)=\sum_{n=0}^{\infty}\cos(n\psi)\mathfrak{P}_{n,p}(\cosh\eta) + (\eta-\log 2)\sinh^p\eta\sum_{n=0}^{p}\epsilon_n\cos(n\psi)\frac{(-p)_n(p-n)!}{(p+n)!}P_p^n(\coth\eta).$$

Furthermore, if we define $\mathfrak{Q}_{n,p}:(1,\infty)\to\mathbf{R}$ as

$$\mathfrak{Q}_{n,p}(\cosh\eta):=\mathfrak{P}_{n,p}(\cosh\eta)+\frac{\epsilon_n(-p)_n(p-n)!}{(p+n)!}(\eta-\log 2)\sinh^p\eta P_p^n(\coth\eta),$$

then we can express g_χ as

$$(\cosh\eta-\cos\psi)^p\log(\cosh\eta-\cos\psi)=\sum_{n=0}^{\infty}\cos(n\psi)\mathfrak{Q}_{n,p}(\cosh\eta). \tag{23}$$

4. Limit derivative approach to the logarithmic Fourier series

We now use a second approach to compute the Fourier series for a logarithmic fundamental solution of the polyharmonic equation (6). We would like to match our results to the computations in §3, which clearly demonstrate different behaviors for the two regimes, $0\leq n\leq p$ and $n\geq p+1$. By applying the identity

$$(\cosh\eta-\cos\psi)^p\log(\cosh\eta-\cos\psi)=\lim_{\nu\to 0}\frac{\partial}{\partial\nu}(\cosh\eta-\cos\psi)^{\nu+p}, \tag{24}$$

where $p\in\mathbf{N}_0$, to

$$(\cosh\eta-\cos\psi)^\nu=\sinh^\nu\eta\sum_{n=0}^{\infty}\frac{(-1)^n\epsilon_n\cos(n\psi)}{(\nu+1)_n}P_\nu^n(\coth\eta), \tag{25}$$

where $\nu\in\mathbf{C}\setminus -\mathbf{N}$ (cf. (3.11b) in Cohl & Dominici (2010) [4]), one can compute the Fourier cosine series of $(\cosh\eta-\cos\psi)^p\log(\cosh\eta-\cos\psi)$, provided availability of the necessary parameter derivatives.

Applying (24) to (25), we obtain

$$
\begin{aligned}
&(\cosh\eta-\cos\psi)^p\log(\cosh\eta-\cos\psi)\\
&\quad=\left[\lim_{\nu\to0}\frac{\partial}{\partial\nu}\sinh^{\nu+p}\eta\right]\sum_{n=0}^{\infty}\frac{(-1)^n\epsilon_n\cos(n\psi)}{(p+1)_n}P_p^n(\coth\eta)\\
&\quad+\sinh^p\eta\sum_{n=0}^{\infty}(-1)^n\epsilon_n\cos(n\psi)\left[\lim_{\nu\to0}\frac{\partial}{\partial\nu}\frac{1}{(\nu+p+1)_n}\right]P_p^n(\coth\eta)\\
&\quad+\sinh^p\eta\sum_{n=0}^{\infty}\frac{(-1)^n\epsilon_n\cos(n\psi)}{(p+1)_n}\left[\lim_{\nu\to0}\frac{\partial}{\partial\nu}P_{\nu+p}^n(\coth\eta)\right].
\end{aligned}
$$

Note that for $p\in\mathbf{Z}$ and $n\in\mathbf{N}_0$, the associated Legendre function of the first kind P_p^n vanishes if $n\geq p+1$. This is easily seen using the Rodrigues-type formula (cf. (14.7.11) in Olver *et al.* (2010) [15])

$$P_p^n(z)=(z^2-1)^{n/2}\frac{d^nP_p(z)}{dz^n},$$

and the fact that $P_p(z)$ is a polynomial in z of degree p. The derivatives are given as follows:

$$\lim_{\nu\to0}\frac{\partial}{\partial\nu}\sinh^{\nu+p}\eta=\sinh^p\eta\log\sinh\eta, \tag{26}$$

$$\lim_{\nu\to0}\frac{\partial}{\partial\nu}\frac{1}{(\nu+p+1)_n}=\frac{p!}{(p+n)!}\left[\psi(p+1)-\psi(p+1+n)\right], \tag{27}$$

and

$$\lim_{\nu\to0}\frac{\partial}{\partial\nu}P_{\nu+p}^n(\coth\eta)=\left[\frac{\partial}{\partial\nu}P_\nu^n(\coth\eta)\right]_{\nu=p}, \tag{28}$$

where $\psi:\mathbf{C}\setminus-\mathbf{N}_0\to\mathbf{C}$ is the digamma function defined in terms of the derivative of the gamma function

$$\frac{d}{dz}\Gamma(z)=:\psi(z)\Gamma(z)$$

(see for instance (5.2.2) in Olver *et al.* (2010) [15]). The degree-derivative of the associated Legendre function of the first kind in (28) is determined using (A2) and

(A3). By collecting terms and using (26), (27), and (28), we obtain

$$
\begin{aligned}
(\cosh\eta-\cos\psi)^p\log(\cosh\eta-\cos\psi) &= (\eta-\log 2)(\cosh\eta-\cos\psi)^p \\
&+p!\sinh^p\eta\sum_{n=0}^{p}\frac{(-1)^n\epsilon_n\cos(n\psi)}{(p+n)!} \\
&\quad\times\left[2\psi(2p+1)-\psi(p+1+n)-\psi(p+1-n)\right]P_p^n(\coth\eta) \\
&+(-1)^p p!\sinh^p\eta\sum_{n=0}^{p-1}\frac{\epsilon_n\cos(n\psi)}{(p+n)!} \\
&\quad\times\sum_{k=0}^{p-n-1}\frac{(-1)^k(2n+2k+1)\left[1+\frac{k!(p+n)!}{(2n+k)!(p-n)!}\right]}{(p-n-k)(p+n+k+1)}P_{n+k}^n(\coth\eta) \\
&+2(-1)^p p!\sinh^p\eta\sum_{n=1}^{p}\frac{(-1)^n\cos(n\psi)}{(p-n)!} \\
&\quad\times\sum_{k=0}^{n-1}\frac{(-1)^k(2k+1)}{(p-k)(p+k+1)}P_k^{-n}(\coth\eta) \\
&+2(-1)^{p+1}p!\sinh^p\eta\sum_{n=p+1}^{\infty}\cos(n\psi)(n-p-1)!P_p^{-n}(\coth\eta). \qquad (29)
\end{aligned}
$$

5. Comparison of the two approaches

The limit derivative approach presented in §4 might be considered, of the two methods, preferred for computing the azimuthal Fourier series for a logarithmic fundamental solution of the polyharmonic equation. This is because it produces azimuthal Fourier coefficients in terms of the well-known special functions, associated Legendre functions. On the other hand, the algebraic approach presented in §3 produces results in terms of the two-parameter logarithmic polynomials $R_p^k(x)$. As far as the author is aware, these polynomials are previously unencountered in the literature. By comparison of the two approaches we see how the logarithmic polynomials $R_p^k(x)$ (potentially a new type of *special function*) are intimately related to the associated Legendre functions. In this section we make this comparison concrete. We should also mention that the following comparison equations resolve to become quite complicated as p increases, and they have been checked for $0\leq p\leq 10$ using Mathematica with the assistance of an algorithm generated using (B2) and (B3) from Appendix B.

By equating the Fourier coefficients using the two approaches we can obtain summation formulae which are satisfied by the logarithmic polynomials. For $n=0$

and $p \geq 1$ we have

$$\sum_{k=1}^{p} \frac{(-1)^{k+1} e^{-k\eta} R_p^k(\cosh\eta)}{k}$$

$$= \sinh^p \eta \left[\psi(2p+1) - \psi(p+1)\right] P_p(\coth\eta)$$

$$+ (-1)^p \sinh^p \eta \sum_{k=0}^{p-1} \frac{(-1)^k (2k+1)}{(p-k)(p+k+1)} P_k(\coth\eta),$$

for $1 \leq n \leq p-1$, $p \geq 2$ we derive

$$\sum_{k=-p}^{n-1} \frac{(-1)^{k+1} e^{k\eta} R_p^k(\cosh\eta)}{n-k} + e^{2n\eta} \sum_{k=-p}^{-n-1} \frac{(-1)^k e^{k\eta} R_p^k(\cosh\eta)}{n+k} = p! e^{n\eta} \sinh^p \eta$$

$$\times \left\{ \frac{(-1)^n}{(p+n)!} \left[2\psi(2p+1) - \psi(p+1+n) - \psi(p+1-n)\right] P_p^n(\coth\eta) \right.$$

$$+ \frac{(-1)^p}{(p+n)!} \sum_{k=0}^{p-n-1} \frac{(-1)^k (2n+2k+1) \left[1 + \frac{k!(p+n)!}{(2n+k)!(p-n)!}\right]}{(p-n-k)(p+n+k+1)} P_{n+k}^n(\coth\eta)$$

$$\left. + \frac{(-1)^{p+n}}{(p-n)!} \sum_{k=0}^{n-1} \frac{(-1)^k (2k+1)}{(p-k)(p+k+1)} P_k^{-n}(\coth\eta) \right\},$$

for $n = p$, $p \geq 1$ we obtain

$$\sum_{k=-p}^{p-1} \frac{(-1)^{k+1} e^{k\eta} R_p^k(\cosh\eta)}{p-k} = p! e^{p\eta} \sinh^p \eta$$

$$\times \left\{ \frac{(-1)^p}{(2p)!} \left[\psi(2p+1) - \psi(1)\right] P_p^p(\coth\eta) \right.$$

$$\left. + \sum_{k=0}^{p-1} \frac{(-1)^k (2k+1)}{(p-k)(p+k+1)} P_k^{-p}(\coth\eta) \right\},$$

and for $n \geq p+1$, $p \geq 0$, we see that

$$\sum_{k=-p}^{p} \frac{(-1)^{k+1} e^{k\eta} R_p^k(\cosh\eta)}{n-k}$$

$$= (-1)^{p+1} p! (n-p-1)! e^{n\eta} \sinh^p \eta P_p^{-n}(\coth\eta). \tag{30}$$

We now have closed-form expressions for the finite terms given by (20), (23) and (20) in terms of associated Legendre functions of the first kind. We also have a proof of the correspondence for the "ending" function $\Re_{n,q-1}$ mentioned in §3. Through (30), the function $\Re_{n,p}$ (cf. (15), (16), (17), (18), and (20)) is directly

related to the associated Legendre function of the first kind, namely

$$\Re_{n,p}(\cosh\eta) = 2(-1)^{p+1}p!(p+n)!e^{n\eta}\sinh^p\eta P_p^{-n}(\coth\eta).$$

Therefore through (10) we have

$$\frac{1}{(\cosh\eta-\cos\psi)^q} = \frac{(-1)^q}{2[(q-1)!]^2\sinh^{2q-1}\eta}\sum_{n=0}^{\infty}\epsilon_n\cos(n\psi)e^{-n\eta}\Re_{n,q-1}(\cosh\eta),$$

which demonstrates the correspondences which was mentioned near (12), (16), and (18) for (20) and (29).

6. Fourier expansion for a logarithmic fundamental solution of the polyharmonic equation

Now that we have computed the Fourier series for g_χ, namely (20) (cf. (23)) and (29), we are in a position to compute the azimuthal Fourier series for a logarithmic fundamental solution of the polyharmonic equation (6).

For instance, using (7), (22) and (29), we have

$$
\begin{aligned}
\mathfrak{l}_k^d(\mathbf{x},\mathbf{x}') &= \frac{1}{2}\left(2RR'\right)^p\left[\log\left(RR'\right)+\eta-\beta_{p,d}\right](\chi^2-1)^{p/2}\\
&\qquad\times\sum_{n=0}^{p}\epsilon_n\cos[n(\phi-\phi')]\frac{(-p)_n(p-n)!}{(p+n)!}P_p^n\left(\frac{\chi}{\sqrt{\chi^2-1}}\right)\\
&+\frac{1}{2}(2RR')^p p!(\chi^2-1)^{p/2}\sum_{n=0}^{p}\frac{(-1)^n\epsilon_n\cos[n(\phi-\phi')]}{(p+n)!}\\
&\qquad\times\left[2\psi(2p+1)-\psi(p+1+n)-\psi(p+1-n)\right]P_p^n\left(\frac{\chi}{\sqrt{\chi^2-1}}\right)\\
&+\frac{1}{2}(2RR')^p p!(\chi^2-1)^{p/2}\sum_{n=0}^{p-1}\frac{\epsilon_n\cos[n(\phi-\phi')]}{(p+n)!}\\
&\qquad\times\sum_{k=0}^{p-n-1}\frac{(-1)^k(2n+2k+1)\left[1+\frac{k!(p+n)!}{(2n+k)!(p-n)!}\right]}{(p-n-k)(p+n+k+1)}P_{n+k}^n\left(\frac{\chi}{\sqrt{\chi^2-1}}\right)\\
&+(2RR')^p(-1)^p p!(\chi^2-1)^{p/2}\sum_{n=1}^{p}\frac{(-1)^n\cos[n(\phi-\phi')]}{(p-n)!}\\
&\qquad\times\sum_{k=0}^{n-1}\frac{(-1)^k(2k+1)}{(p-k)(p+k+1)}P_k^{-n}\left(\frac{\chi}{\sqrt{\chi^2-1}}\right)\\
&+(2RR')^p(-1)^{p+1}p!(\chi^2-1)^{p/2}\\
&\qquad\times\sum_{n=p+1}^{\infty}\cos[n(\phi-\phi')](n-p-1)!P_p^{-n}\left(\frac{\chi}{\sqrt{\chi^2-1}}\right).
\end{aligned}
$$

Alternatively, using (7), (9), and (21) we obtain

$$\begin{aligned}
\mathfrak{l}_k^d(\mathbf{x},\mathbf{x}') = \frac{1}{2}\left(2RR'\right)^p\left[\log\left(RR'\right)+\eta-\beta_{p,d}\right](\chi^2-1)^{p/2} \\
\times\sum_{n=0}^{p}\epsilon_n\cos[n(\phi-\phi']\frac{(-p)_n(p-n)!}{(p+n)!}P_p^n\left(\frac{\chi}{\sqrt{\chi^2-1}}\right), \\
+(2RR')^p\sum_{n=0}^{p}\cos[n(\phi-\phi')]\left(\chi-\sqrt{\chi^2-1}\right)^n \\
\times\sum_{k=-p}^{n-1}\frac{(-1)^{k+1}\left(\chi+\sqrt{\chi^2-1}\right)^k R_p^k(\chi)}{n-k} \\
-(2RR')^p\sum_{n=1}^{p-1}\cos[n(\phi-\phi')]\left(\chi+\sqrt{\chi^2-1}\right)^n \\
\times\sum_{k=-p}^{-n-1}\frac{(-1)^{k+1}\left(\chi+\sqrt{\chi^2-1}\right)^k R_p^k(\chi)}{n+k} \\
+(2RR')^p\sum_{n=p+1}^{\infty}\cos[n(\phi-\phi')]\left(\chi-\sqrt{\chi^2-1}\right)^n \\
\times\sum_{k=-p}^{p}\frac{(-1)^{k+1}\left(\chi+\sqrt{\chi^2-1}\right)^k R_p^k(\chi)}{n-k}.
\end{aligned}$$

Using these formulae, we can, for instance, obtain the axisymmetric component of a logarithmic fundamental solution of the polyharmonic equation, namely

$$\begin{aligned}
\mathfrak{l}_k^d(\mathbf{x},\mathbf{x}')\Big|_{n=0} = \frac{1}{2}(2RR')^p(\chi^2-1)^{p/2} \\
\times\left[\log(RR')+\eta-\beta_{p,d}+2\psi(2p+1)-2\psi(p+1)\right]P_p\left(\frac{\chi}{\sqrt{\chi^2-1}}\right) \\
+(2RR')^p(\chi^2-1)^{p/2}\sum_{k=0}^{p-1}\frac{(-1)^k(2k+1)}{(p-k)(p+k+1)}P_k\left(\frac{\chi}{\sqrt{\chi^2-1}}\right),
\end{aligned}$$

or

$$\begin{aligned}
\mathfrak{l}_k^d(\mathbf{x},\mathbf{x}')\Big|_{n=0} = \frac{1}{2}(2RR')^p(\chi^2-1)^{p/2}\left[\log(RR')+\eta-\beta_{p,d}\right]P_p\left(\frac{\chi}{\sqrt{\chi^2-1}}\right) \\
+(2RR')^p\sum_{k=1}^{p}\frac{(-1)^{k+1}(\chi-\sqrt{\chi^2-1})^k R_p^k(\chi)}{k}.
\end{aligned}$$

The above expressions for the axisymmetric component of a fundamental solution of the polyharmonic equation is one type of expression sought after in Tsai, Chen & Hsu (2009) [19].

Appendix A. Derivatives with respect to the degree of certain integer-order associated Legendre functions of the first kind

The derivative with respect to its degree for the associated Legendre function of the first kind evaluated at the zero degree is given in §4.4.3 of Magnus, Oberhettinger & Soni (1966) [11] as

$$\left[\frac{\partial}{\partial\nu}P_\nu(z)\right]_{\nu=0} = \frac{z-1}{2}{}_2F_1\left(1,1;2;\frac{1-z}{2}\right). \tag{A1}$$

An important generalization of this formula has recently been derived in Szmytkowski (2011) [18]. The degree-derivative of the associated Legendre function of the first kind for $p,m\in\mathbf{N}_0$ and $0\le m\le p$ (cf. (5.12) in Szmytkowski (2011) [18]) is given by

$$\begin{aligned}\left[\frac{\partial}{\partial\nu}P_\nu^m(z)\right]_{\nu=p} &= P_p^m(z)\log\frac{z+1}{2}\\ &+ \left[2\psi(2p+1)-\psi(p+1)-\psi(p-m+1)\right]P_p^m(z)\\ &+ (-1)^{p+m}\sum_{k=0}^{p-m-1}(-1)^k\frac{(2k+2m+1)\left[1+\frac{k!(p+m)!}{(k+2m)!(p-m)!}\right]}{(p-m-k)(p+m+k+1)}P_{k+m}^m(z)\\ &+ (-1)^p\frac{(p+m)!}{(p-m)!}\sum_{k=0}^{m-1}(-1)^k\frac{2k+1}{(p-k)(p+k+1)}P_k^{-m}(z),\end{aligned} \tag{A2}$$

and for $m\ge p+1$ (cf. (5.16) in Szmytkowski (2011) [18]) there is

$$\left[\frac{\partial}{\partial\nu}P_\nu^m(z)\right]_{\nu=p} = (-1)^{p+n+1}(p+n)!(n-p-1)!P_p^{-n}(z). \tag{A3}$$

Some special cases of (A2) include for $m=0$

$$\begin{aligned}\left[\frac{\partial}{\partial\nu}P_\nu(z)\right]_{\nu=p} &= P_p(z)\log\frac{z+1}{2}+2\left[\psi(2p+1)-\psi(p+1)\right]P_p(z)\\ &+ 2(-1)^p\sum_{k=0}^{p-1}(-1)^k\frac{2k+1}{(p-k)(p+k+1)}P_k(z),\end{aligned}$$

for $m = p$

$$\left[\frac{\partial}{\partial\nu}P_\nu^p(z)\right]_{\nu=p} = P_p^p(z)\log\frac{z+1}{2} + \left[2\psi(2p+1) - \psi(p+1) + \gamma\right]P_p^p(z)$$

$$+ (-1)^p(2p)!\sum_{k=0}^{p-1}(-1)^k\frac{2k+1}{(p-k)(p+k+1)}P_k^{-p}(z),$$

where $\gamma = -\psi(1)$ is Euler's constant $\gamma \approx 0.57721566490$. Of course we also have for $m = p = 0$

$$\left[\frac{\partial}{\partial\nu}P_\nu(z)\right]_{\nu=0} = \log\frac{z+1}{2},$$

which exactly matches (A1).

Appendix B. The logarithmic polynomials

The logarithmic polynomials R_p^k are nonvanishing only for $-p \leq k \leq p$ and by construction, they are satisfied by the following recurrence relation

$$R_p^k(x) = \frac{1}{2}R_{p-1}^{k-1}(x) + xR_{p-1}^k(x) + \frac{1}{2}R_{p-1}^{k+1}(x). \tag{B1}$$

From (11) we have that $R_0^0(x) = 1$. This gives us the starting point for the recursion. We conjecture that the derivative of the logarithmic polynomials R_p^k is given by

$$\frac{d}{dx}R_p^{\pm k}(x) = pR_{p-1}^{\pm k}(x).$$

It is evident by construction that these polynomials are even in the index k, i.e.,

$$R_p^k(x) = R_p^{-k}(x).$$

Some of the first few logarithmic polynomials are given by

$$R_0^0(x) = 1,$$

$$R_1^0(x) = x,\ R_1^{\pm1}(x) = \frac{1}{2}$$

$$R_2^0(x) = \frac{1}{2} + x^2,\ R_2^{\pm1}(x) = x,\ R_2^{\pm2}(x) = \frac{1}{4},$$

$$R_3^0(x) = \frac{3}{2}x + x^3,\ R_3^{\pm1}(x) = \frac{3}{8} + \frac{3}{2}x^2,\ R_3^{\pm2}(x) = \frac{3}{4}x,\ R_3^{\pm3}(x) = \frac{1}{8}.$$

We can find the generating function for the logarithmic polynomials as follows. Let

$$F(x,y,z) = \sum_{p=0}^{\infty}\sum_{k=-\infty}^{\infty}R_p^k(x)y^kz^p$$

be the generating function for the logarithmic polynomials R_p^k. If we define the function

$$S_p(x,y) = \sum_{k=-\infty}^{\infty} R_p^k(x) y^k,$$

then using the recurrence relation for R_p^k (B1) we can show

$$S_p(x,y) = \left(x + \frac{1}{2}\left(y + \frac{1}{y}\right)\right) S_{p-1}(x,y).$$

Combining this result along with the fact that $R_0^0(x) = 1$, we have

$$S_p(x,y) = \left(x + \frac{1}{2}\left(y + \frac{1}{y}\right)\right)^p,$$

so therefore the generating function for the logarithmic polynomials R_p^k is given by

$$F(x,y,z) = \frac{1}{1 - z\left(x + \frac{1}{2}\left(y + \frac{1}{y}\right)\right)}.$$

An algorithm for generating the logarithmic polynomials can be obtained by solving the following set of difference equations

$$\left.\begin{aligned} a_0(p) &= \tfrac{1}{2} a_0(p-1) \\ a_1(p) &= \tfrac{1}{2} a_1(p-1) + x a_0(p-1) \\ a_2(p) &= \tfrac{1}{2} a_2(p-1) + x a_1(p-1) + \tfrac{1}{2} a_0(p-1) \\ &\vdots \\ a_n(p) &= \tfrac{1}{2} a_n(p-1) + x a_{n-1}(p-1) + \tfrac{1}{2} a_{n-2}(p-1) \end{aligned}\right\}, \tag{B2}$$

subject to the boundary conditions

$$\left.\begin{aligned} a_0(0) &= 1 \\ a_1(1) &= x a_0(0) \\ a_2(2) &= x a_1(1) + a_0(1) \\ &\vdots \\ a_n(n) &= x a_{n-1}(n-1) + a_{n-2}(n-1) \end{aligned}\right\}, \tag{B3}$$

where $R_p^k = a_{p-|k|}(p)$ is given along diagonals for a fixed $p - |k|$. For instance, one can obtain $R_p^{\pm p}(x) = \frac{1}{2^p}$, for $p \geq 0$, $R_p^{\pm(p-1)}(x) = \frac{p}{2^{p-1}} x$, for $p \geq 1$ and $R_p^{\pm(p-2)}(x) = \frac{p}{2^p} + \frac{p(p-1)}{2^{p-1}} x^2$, for $p \geq 2$, etc.

Acknowledgements

I would like to thank Tom ter Elst, Matthew Auger, and Radosław Szmytkowski for valuable discussions. I would also like to thank Shenghui Yang at Wolfram Research for valuable assistance in generating an algorithm to symbolically compute the logarithmic polynomials. I acknowledge funding for time to write this paper from the Dean of the Faculty of Science at the University of Auckland in the form of a three month stipend to enhance University of Auckland 2012 PBRF Performance. Part of this work was conducted while H. S. Cohl was a National Research Council Research Postdoctoral Associate in the Information Technology Laboratory at the National Institute of Standards and Technology, Gaithersburg, Maryland, U.S.A.

References

[1] M. Abramowitz and I. A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. U.S. Government Printing Office, Washington, D.C., 1972.

[2] J. B. Boyling. Green's functions for polynomials in the Laplacian. *Zeitschrift für Angewandte Mathematik und Physik*, 47(3):485–492, 1996.

[3] H. S. Cohl. *Fourier and Gegenbauer expansions for fundamental solutions of the Laplacian and powers in* $\mathbf{R}^d$ *and* $\mathbf{H}^d$. PhD thesis, The University of Auckland, 2010. xiv+190 pages.

[4] H. S. Cohl and D. E. Dominici. Generalized Heine's identity for complex Fourier series of binomials. *Proceedings of the Royal Society A*, 467:333–345, 2010.

[5] H. S. Cohl and J. E. Tohline. A Compact Cylindrical Green's Function Expansion for the Solution of Potential Problems. *The Astrophysical Journal*, 527:86–101, December 1999.

[6] I. M. Gel'fand and G. E. Shilov. *Generalized functions. Vol. 1.* Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1964 [1977]. Properties and operations, Translated from the Russian by Eugene Saletan.

[7] BT Helenbrook. Mesh deformation using the biharmonic operator. *International Journal for Numerical Methods in Engineering*, 56(7):1007–1021, 2003.

[8] B. J. Kirby. *Micro- and Nanoscale Fluid Mechanics: Transport in Microfluidic Devices*. Cambridge University Press, Cambridge, 2010.

[9] W.M. Lai, D. Rubin, and E. Krempl. *Introduction to continuum mechanics, revised edition, SI/metric units*. Pergamon, Oxford, UK, 1978.

[10] S. A. Lurie and V. V. Vasiliev. *The biharmonic problem in the theory of elasticity*. Gordon and Breach Publishers, Luxembourg, 1995.

[11] W. Magnus, F. Oberhettinger, and R. P. Soni. *Formulas and theorems for the special functions of mathematical physics*. Third enlarged edition. Die Grundlehren der mathematischen Wissenschaften, Band 52. Springer-Verlag New York, Inc., New York, 1966.

[12] W. Miller, Jr. *Symmetry and separation of variables*. Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1977. With a foreword by Richard Askey, Encyclopedia of Mathematics and its Applications, Vol. 4.

[13] J. Monterde and H. Ugail. On harmonic and biharmonic Bézier surfaces. *Computer Aided Geometric Design*, 21(7):697–715, 2004.

[14] M. Nicolescu. *Les Fonctions Polyharmoniques*. Hermann, Paris, 1936.

[15] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, editors. *NIST handbook of mathematical functions*. Cambridge University Press, Cambridge, 2010.

[16] L. Schwartz. *Théorie des distributions. Tome I.* Actualités Sci. Ind., no. 1091 = Publ. Inst. Math. Univ. Strasbourg 9. Hermann, Paris, 1950.

[17] S. L. Sobolev. *Cubature formulas and modern analysis: an introduction*. Oxonian, New Delhi, 1992.

[18] R. Szmytkowski. On the derivative of the associated Legendre function of the first kind of integer order with respect to its degree (with applications to the construction of the associated Legendre function of the second kind of integer degree and order). *Journal of Mathematical Chemistry*, 49(7):1436–1477, June 2011.

[19] C. S. Tsai, C.-C. and Chen and T.-W. Hsu. The method of particular solutions for solving axisymmetric polyharmonic and poly-Helmholtz equations. *Engineering Analysis with Boundary Elements*, 33(12):1396–1402, 2009.

[20] H. Ugail. *Partial Differential Equations for Geometric Design*. Springer-Verlag, London, 2011.

[21] D. Vautherin. Hartree-Fock Calculations with Skyrme's Interaction. II. Axially Deformed Nuclei. *Physical Review C*, 7:296–316, 1973.